



Research Article

The work is licensed under



Modulations in Haematological Aspects of Wistar Rats Exposed to Sublethal Doses of Fipronil under Subchronic Duration

Kartheek RM, David M*

Environmental Toxicology and Molecular Biology Laboratory, Department of PG Studies and Research in Zoology, Karnatak University, Dharwad- 580003

*Corresponding Author: David M, Environmental Toxicology and Molecular Biology Laboratory, Department of PG Studies and Research in Zoology, Karnatak University, Dharwad- 580003

Received: 25 August 2017 Revised: 09 September 2017 Accepted: 11 September 2017

ABSTRACT

Fipronil (FPN) a phenylpyrazole insecticide is one of the preferred agricultural chemical known for controlling insect pests. Assessment of its safe and toxic levels has become a matter of serious concern as it is occupationally exposed to the farming community. The current investigation was thus aimed to elucidate the changes in haematological indices of Wistar rats on exposure to FPN. Healthy and mature male Wistar rats were distributed into four groups, namely C, E1, E2 and E3 and received 0.0, 6.46, 12.12 and 32.33mg/kg body weight of FPN respectively for a duration of 90 days. The results obtained indicated significant ($*p < 0.05$) changes in RBC, WBC, Hb, PCV, MCH and MCHC values unlike MCV as compared to control. The decline noticed in RBC suggested anaemic conditions of rats. Significant ($*p < 0.05$) elevation in WBC at E1 and subsequent decline at E2 and E3 was also witnessed under current study. As the modulatory outcome of FPN is evident on haematological aspects, it is thereby suggested that FPN under selected doses should be avoided for application into agricultural fields unless precautionary measures are taken to avoid exposure of toxicant as it threatens the health of the mammalian class resulting in compromised haematological condition.

Keyword: Fipronil; haematotoxicity; immune system; red blood corpuscles; Wistar rats

INTRODUCTION

The widespread use of pesticides in agriculture can lead to soil, water and air contamination resulting in adverse effects on inhabiting non-target organisms [1]. A number of pesticides are presented mainly into the agricultural landscapes as racemates even though their activity usually results from a preferential reactivity of only one enantiomer [2]. The pesticides currently in use include a wide variety of chemicals with great differences in their mode of action, uptake by the body, metabolism and elimination from the body, and causes the

adverse health conditions to the non-target organisms including humans [3]. In the present era a large diverse group of insecticides are utilized against the insect pests thereby enhancing agriculture yield, but they eventually are found to reach the aquatic ecosystem through agriculture runoff ultimately entering the food chain in more adequate amounts [4]. After organochlorines [5], carbamates [6], synthetic pyrethroids [7] and organophosphates [8]; new class insecticides are being explored for their ability to selectively kill insect pests with lesser or no damage to non-target organisms.

Increased attention is being given to the mammalian toxicity studies as the adverse effects caused by new and conventional group of pesticides are innumerable [9]. Toxicity of pesticides on mammalian species include effect on biochemical, neurological, and haematological factors [10, 11].

Haematological indices are the important biomarker for suggesting altered internal and/or external environment of animals and variations in their indices within an individual can result in inadequate responses to chemical stressors; however, these variations are nonspecific to a wide range of substances [12]. Exposure to chemical pollutants especially pesticides is thought to either increase or decrease in hematological levels [13] often affecting the survival of exposed animal. Hence, in the evaluation of health conditions and outcome from the pesticide induced toxicity within organisms, parameters under hematological and biochemical aspects are exercised as indicators [14-16]. Even though broad attempts of studies have been performed to demarcate the haematotoxic potentials of conventional group of pesticides, the same for new class insecticides like phenylpyrazole have been understudied for their ability to cause disturbances in blood of an organism.

Fipronil (5- amino- 1- (2, 6-dichloro- α , α , α -trifluoro- p- tolyl)- 4-trifluoromethylsulfinylpyrazole-3- carbonitrile) (FPN) is broad spectrum, first N- phenylpyrazole insecticide introduced with an objective of controlling insect pest [17]. It belongs to the group of "new generation" insecticide as its mode of action is dissimilar to pyrethroids (sodium channel blockers), organophosphates, and carbamate (cholinesterase inhibitors) to which a variety of insect pests have gained resistance [18]. Nonetheless, the mechanism of action of FPN has been previously discussed by Cole *et al.* [19], who found that the phenylpyrazole insecticide interferes with the γ -aminobutyric acid (GABA) gated channel thereby disrupting normal nerve influx transmission and at sufficient dose causes excessive neural excitation, severe paralysis and finally death [18, 20, 21]. Even though FPN demonstrates a selective toxicity by having a higher binding affinity toward the GABA-regulated chloride channels of insects (LC_{50} can be low as 24.8 nM or \sim 11.7 μ g/L) than for mammalian GABA

receptors [22], its ability to cause haematological alterations through other modes cannot be ignored. Hence, in the present study, an attempt has been made to evaluate the impact of FPN on blood indices and determine safe and toxic dose levels against wistar rats following exposure to subchronic duration.

MATERIALS AND METHODS

Animal procurement

Male Wistar albino rats (6 weeks) were obtained from animal house facility, Department of PG studies and Research in Zoology, Karnatak University, Dharwad, Karnataka, India. Rats were housed in the polypropylene cages with *ad libitum* access to standard pellet feed and drinking water. The room was maintained under a 12/12 h light– dark cycle, an ambient temperature of 23-30 °C with a relative humidity of 45 (\pm 15)%.

Experimental design and test doses

For a 90-day study of oral toxicity, the 6 week old Wistar rats were randomly assigned into four groups (C, E1, E2 and E3) of six males each. Group C of untreated rats served as the control group. Rats in the group E1, E2 and E3 were exposed to of 0.0, 6.46, 12.12 and 32.33mg/kg body weight of FPN dose in drinking water.

Haematological indices

Blood was drawn by puncturing the retro-orbital plexus under diethyl ether anesthesia. Whole blood for haematogram was collected in vials containing the anticoagulant, ethylene diamine tetra-acetic acid. Total erythrocytes (RBC), total leukocyte count (WBC), packed cell volume (PCV), Mean corpuscular volume (MCV), MCH and MCHC were estimated by using the standard procedure of Rodak [23], Hemoglobin (Hb) by the method of Van Kampen and Zijlstra, [24].

Statistical analysis

The values for haematological indices are reported as the mean \pm standard error of the mean (SEM) obtained from triplicates. The data were subjected to one-way analysis of variance and further subjected to Tukey's test for post hoc analysis by defining the significance level at * p <0.05.

Ethical committee

All rats were housed for a duration of 1 week for acclimatization before initiation of the experiments. The maintenance of experimental rats and all the procedures implemented are in accordance with standard guidelines issued by CPCSEA followed with approval of the Institutional Animal Ethics Committee (IAEC) of the institute.

RESULTS AND DISCUSSION

Changes in haematological indices are considered to be one of the crucial biomarkers of physiological stress [25]. The present investigation indicated significant ($p < 0.05$) variations in haematological indices of rats exposed to different sublethal doses of FPN for 90 days and have been presented in figure 1-7. No changes were observed in control rats which advertised normal count. The rats under FPN stress however advertised changes in RBC faction with percent decline of -19.57, -19.01 and -57.60% for E1, E2 and E3 respectively. Changes in RBC count which was found to be in an irregular trend, which suggests the involvement of FPN in either demolishing the RBC concentrates or intervening in erythropoietic mechanism as reported previously by Mehra et al. [26]. Since the RBC's account for transportation of oxygen and deoxygenated blood, variation in this ratio within the individual is thought to play a crucial role in maintenance of balance between oxygen and carbon dioxide levels [27]. Our reports are in agreement with Magar and Dube [28], Goel et al. [29] and Ambali et al. [30] who stated that exposure to toxicants under long term durations have been known to cause anomalies including anemia in non-target organisms [31]. Changes in WBC count was found to be in irregular pattern with an elevated cell count at E1 (41.57%) and E2 (28.09%) unlike for E3 wherein a decline of -57.30% was witnessed. The elevation in WBC may be due to the activation of the animal's defence mechanism against the toxic invasion by FPN concentrates. However, the E3 group demonstrated significant decline in WBC's which indicates the probabilities of systemic damage to the immune system due to persistence of high doses of FPN. Similar reports have been given by Lohner et al. [32] and Shahi and Singh, [33] who reported decline and increase in WBC's respectively under different

conditions. Also Janeway et al. [34], explained that decreased number of lymphocytes may be due to reduced production or rapid removal from circulation and subsequent destruction. The reduction of lymphocytes is indicative of immunosuppressive effects and has been previously reported for organophosphates. Moreover, the results obtained by Goel et al. [29] confirmed that, decrease in leucocyte counts following intoxication with another organophosphate, Chlorpyrifos, could be attributed either to the slower rate of production of leucocytes or due to their inhibited release into the blood circulation. It is to be noted that the report on WBC decline is reported for the first time against the FPN for the subchronic duration of 90 days. Haemoglobin is the iron-containing oxygen-transport metallo-protein in the red blood cells of all vertebrates [35].

Haemoglobin displays the function of oxygen transportation within the tissues of the animals for oxidation of ingested food so as to release energy for the other body functions as well as transport carbon dioxide out of the body of animals. Changes in were noted for Hb which advertised highest decline at E3 (-55.82) followed by E2 (-23.66) and E1 (-18.92), which further suggests lesser difference between E2 and E1. The variation in Hb and decline in RBC was found to be dependent on the dose level of FPN administered suggests that FPN induces toxicity against RBC's of rats resulting in increased rate of erythrocytes destruction which could be taking place in hemopoietic organs.

PCV under current investigation demonstrated highest damage under E3 group followed by E1 and E2. However, the change was found to be significant ($p < 0.05$) under all the exposed group. Reports by Isaac et al. [36] have suggested that PCV is primarily associated in the transport of oxygen and absorbed nutrients. Since the present investigation was recorded for decreased PCV levels, the possibility of lack of oxygen and nutrient transportation cannot be overlooked. The reduction in both quality and quantity of PCV may be a consequence of severe haemorrhage and is often correlated with RBC count and Hb levels. This in turn results in the dilution of blood caused by the influx of cells and fluids from body stores [37]. In fact, Hb concentration and PCV values are directly correlating with RBC's count, this part of our study is in support of reports from El-Bakary et

al. [38], and the mechanism behind could be mainly due to the synergistic link among these blood parameters in all vertebrates. The association between the total RBC count and the

corresponding haemoglobin levels has been often compared with the haematocrit value and has been reported for other vertebrates including humans [39].




Fig. 1: Changes in total erythrocyte count of wistar rats following exposure to different sublethal doses of fipronil for 90 days




Fig. 2: Changes in total leucocyte count of wistar rats following exposure to different sublethal doses of fipronil for 90 days




Fig. 3: Changes in haemoglobin level of wistar rats following exposure to different sublethal doses of fipronil for 90 days




Fig. 4: Changes in levels of packed cell volume of wistar rats following exposure to different sublethal doses of fipronil for 90 days

Change in MCV under all the groups could be as a consequence of varying erythrocyte size that in turn may be due to the process of osmotic imbalance in fluid conditions. Our results are in agreement with reports of Al-Amoudi, [40], who reported similar trend in blood indices of rats after exposure to organophosphate. The data for MCV unlike for MCH and MCHC advertised insignificant variation. A slight decline was observed at E1 (-0.71%) and a trivial increase

was noted for E2 (0.21%). However, E3 showed a significant elevation of 46.89%. Evaluation of MCV thus provides an information on the size and status of erythrocytes within the exposed organism [41]. Reports given by Anandkumar et al.[42] suggested increase in MCV that is caused due to endosmosis. The haemodilution is the consequence of endosmosis that leads to the passage of solvent from less concentrated solution to more concentrated one and has been

found to increase the MCV value. Previously Garg, *et al.* [43], Dhanapakiam and Ramasamy,

[44] reported increase in MCV within non-target organisms exposed to toxicants.




Fig. 5: Changes in levels of mean corpuscular volume of wistar rats following exposure to different sublethal doses of fipronil for 90 days




Fig. 6: Changes in levels of mean corpuscular haemoglobin of wistar rats following exposure to different sublethal doses of fipronil for 90 days




Fig.7: Changes in levels of mean corpuscular haemoglobin concentration of wistar rats following exposure to different sublethal doses of fipronil for 90 days

Under MCH, the values suggested significant variation ($p < 0.05$) with percent change of 97.3, 84.5, 103.9 % for E1, E2 and E3 respectively. The swelling up of RBC has been through determination of MCHC which has been considered as a critical indicator under haematological studies [45]. The results from the present study speculates the possibility of FPN inducing physical change in rat erythrocyte morphology from the normal discoid shape to other forms including echinocytes, dacrocytes

and schistocytes. Echinocytes transformation plays an anti-haemolytic role – the expansion of the plasma membrane increases the cell-volume ratio/membrane-area, thus allowing swelling of the cells before lysing [46]. The current investigation revealed MCHC values which were found to be significant ($p < 0.05$) and elevated under the FPN intake resulting in a percent change of 98.86, 84.21 and 38.93% for E1, E2 and E3 respectively. As MCH and MCHC are derived from Hb and RBC, any sort of alteration

in the levels of Hb and RBC would commonly result in the alteration of MCH and MCHC. Atamanalp, *et al.* [47] reported a marked rise in the values of MCHC, MCH and MCV in *O. mykiss* on exposure cypermethrin. Changes in MCV, MCH and MCHC levels has been reported by Parma, *et al.* [48], in a ray-finned fish *Prochilodus lineatus* upon exposure to 0.6 ppm of cypermethrin which suggested the haematotoxic potential. As reported previously, various environmental toxicants are known to induce changes in organs like brain [49], liver [50, 51], kidney [52] and spleen [53] causing impairments, resulting in the persistence of FPN residues in blood thereby causing significant variations in its indices. The current work is viewed as a nutshell and further needs molecular investigations to elucidate the mechanism of FPN in demolition of haematological integrity of the exposed organism.

CONCLUSION

FPN exposure to rats was found to alter total RBC and WBC counts suggesting anaemic condition and instigation of immune response respectively. The decline of Hb as observed under current investigation indicates the possibilities of hypoxic conditions of exposed animals. Significant change in PCV, MCH and MCHC indicates the direct impact on overall haematological aspects within the exposed animal. Thus based on the outcome, it could be ascertained that FPN is toxic at all the selected sublethal doses when exposed for a duration of 90 days. Hence it is suggested that care is to be taken when the toxicant is used or sprayed under mammalian vicinity.

ACKNOWLEDGEMENTS

The authors are thankful to the University Grants Commission for providing financial assistance through UGC SAP scheme [No. F.4-18/2015/DSA-I (SAP-II)] and UGC-UPE/FAR I [(37-245/2009 (SR)]. The authors are also thankful to the Department of Science and Technology for providing financial support through DST PURSE Phase II Program.

CONFLICT OF INTEREST

The authors confirmed that there is no conflict of interest for this research paper.

REFERENCES

1. Karthek RM, David M. Fipronil induced modulations in biochemical and histopathological aspects of male Wistar albino rats: A subchronic study. *J Environ Bios* 2016; 5:26-32.
2. Nillos MG, Gan J and Schlenk D. Chirality of organophosphorus pesticides: analysis and toxicity *J Chromatogr B* 2010; 878:1277-1284.
3. Damalas CA, Koutroubas SD. Farmers' Exposure to Pesticides: Toxicity Types and Ways of Prevention *Toxics* 2016; 4:1-10.
4. David M, Karthek RM. Malathion acute toxicity in tadpoles of *Duttaphrynus melanostictus*, morphological and behavioural study. *The J Bas Appl Zool* 2015;72:1-7
5. Singh Z, Kaur J, Kaur R, Hundal SS. Toxic Effects of Organochlorine Pesticides: A Review. *Amer J Bios* 2016; 4:11-18.
6. Cable GG, Doherty S. Acute carbamate and organochlorine toxicity causing convulsions in an agricultural pilot: a case report. *Aviat Space Environ Med* 1999; 70: 68-72.
7. David M, Shrinivas J, Karthek RM. Biochemical modulations in *Duttaphrynus melanostictus* tadpoles, following exposure to commercial formulations of cypermethrin: An overlooked impact of extensive cypermethrin use. *J Appl Biol Biotechnol* 2016; 4:32-37.
8. Sarabia L, Maurer I, Bustos-Obregón E. Melatonin prevents damage elicited by the organophosphorous pesticide diazinon on the mouse testis. *Ecotoxicol Environ Saf* 2009; 72: 938-942.
9. David M, Karthek RM. Sodium cyanide induced histopathological changes in kidney of fresh water fish *Cyprinus carpio* under sublethal exposure. *Int J Pharm Chem Biol Sci* 2014; 4:634-639.
10. Mushigeri SB, David M. Accumulation of fenvalerate and related changes in lactate and succinate dehydrogenases activity in functionally different tissues of the freshwater fish, *Cirrhinus mrigala* (Hamilton). *J Bas Clin Physiol Pharmacol* 2004; 15:143-152.
11. David M, Karthek RM. Sodium cyanide induced biochemical and histopathological changes in fresh water fish *Cyprinus carpio*

- under sublethal exposure. *Int J Toxicol Appl Pharmacol* 2014; 4:64-69.
12. Prashanth MS and David M. Changes in nitrogen metabolism of the freshwater fish *Cirrhinus mrigala* following exposure to cypermethrin. *J Basic Clin Physiol Pharmacol* 2006; 17:63-70
 13. Kori-Siakpere O, EC Oboh. Haematological effects of sublethal concentrations of tobacco leaf dust on the African catfish: *Clarias gariepinus* (Burchell 1822). *Arch Appl Sci Res* 2011; 3: 93-502.
 14. Thrall MA. Haematology of amphibians. Veterinary haematology and clinical chemistry: Text and clinical case presentations. Philadelphia: Lippincott Williams and Wilkins; 2004.
 15. Pimpao CT, Zampronio AR, De Assis HCS. Effects of deltamethrin on hematological parameters and enzymatic activity in *Ancistrus multispinis* (Pisces, Teleostei). *Pest Biochem Physiol* 2007; 88:122-127.
 16. Rao JV. Biochemical alterations in euryhaline fish *Oreochromis mossambicus* exposed to sub-lethal concentrations of an organophosphorus insecticide, monocrotophos. *Chemosph* 2006; 65:1814-1820.
 17. Simon-Delso N, Amaral-Rogers V, Belzunces LP, Bonmatin JM, Chagnon M, Downs C et al. Systemic insecticides (neonicotinoids and fipronil): trends, uses, mode of action and metabolites. *Environ Sci Pollut Res* 2015; 22:5-34.
 18. Aajoud A, Ravanel P, Tissut M. Fipronil metabolism and dissipation in a simplified aquatic ecosystem. *J Agric Food Chem* 2003; 51:1347-1352.
 19. Cole LM, Nicholson RA, Casida JE. Action of phenylpyrazole insecticides at the GABA-gated chloride channel. *Pestic Biochem Phys* 1993; 46:47-54.
 20. Bobe A, Cooper JF, Coste CM, Muller MA. Behaviour of fipronil in soil under Sahelian plain field conditions. *Pestic Sci* 1998; 52: 275-281.
 21. Gant DB, Chalmers AE, Wolff MA, Hoffman HB, Bushey DF. Fipronil: action at the GABA receptor. *Rev Toxicol* 1998; 2:147-156.
 22. Hainzl D, Casida JE. Fipronil insecticide: novel photochemical desulfinylation with retention of neurotoxicity. *Proc Natl Acad Sci* 1996; 93:12764-12777.
 23. Rodak LC. Routine testing in haematology In: Rodak BE, eds, *Diagnostic Haematology*, WB Saunders, Philadelphia, London, Toronto; 1995.
 24. Van Kampen E, Zijlstra WG. Standardization of hemoglobinometry. II. The hemiglobincyanide method. *Clin Chim Acta* 1961; 6:538-544.
 25. Akinrotimi OA, Abu OMG, Ansa EJ, Edun OM, George OS. Hematological responses of *Tilapia guineensis* to acute stress. *Int J Nat Appl Sci* 2009; 5:338-348.
 26. Mehra BL, Sharma P, Kaushik U, Joshi SC. Effect of Fytolan on Haematology and Serum Parameters of Male Albino Rats. *Int J Pharma Res Health Sci* 2014; 2: 332-338.
 27. Dzik WH. The air we breathe: three vital respiratory gases and the red blood cell: oxygen, nitric oxide, and carbon dioxide. *Transfus* 2011; 51:676-685.
 28. Magar RS, Dube KV. Impact of malathion on some haematological parameters of *Channa punctatus* (Bloch). *Int J Biomed Adv Res* 2012; 03: 683-685.
 29. Goel A, Danni V, Dhawan DK. Role of zinc in mitigating the toxic effects of chlorpyrifos on hematological alterations and electron microscopic observations in rat blood. *Bio Metals* 2006; 19: 483-492.
 30. Ambali SF, Abubakar AT, Shittu M, Yaqub LS, Anafi SB, Abdullahi A. Chlorpyrifos-induced alterations of hematological parameters in Wistar rats: Ameliorative effect of zinc. *Res J Environ Toxicol* 2010; 4:55-66.
 31. Bigoniyaa P, Sahu T, Tiwari V. Hematological and biochemical effects of sub-chronic artesunate exposure in rats. *Toxicol Rep* 2015; 2: 280-288.
 32. Lohner TW, Reash RJ, Williams M. Assessment of tolerant sunfish populations (*Lepomis* sp.) inhabiting selenium-laden coal ash effluents. Haematological and population level assessment. *Ecotoxicol Environ Saf* 2001; 50: 203-216.
 33. Shahi J, Singh A. Genotoxic and haematological effect of commonly used fungicide on fish *Clarias batracus*. *J Biol Earth Sci* 2014; 4:137-143.
 34. Janeway C, Travers P, Walport M, Shlomchik M. Immunobiology: The Immune System in Health and Disease. New York, NY, USA: Garland Science; 2005.

35. Maton A, Hopkins RLJ, McLaughlin C W, Johnson S, Warner CW, La Hart D, Wright JD. Human Biology and Health. Englewood Cliffs, New Jersey, USA: Prentice Hall; 1993.
36. Isaac LJ, Abah G, Akpan B and Ekaette IU. Haematological properties of different breeds and sexes of rabbits. Proceedings of the 18th Annual Conference of Animal Science Association of Nigeria; 2013, p 24.
37. Sharma RK, Saxena Y. Haematological study of Indian desert gerbil, *Meriones hurrianae* (Jerdon) after brodifacoum poisoning. Indian J Comp Animal Physiol 1983;1:23-28.
38. El-Bakary AS, Abdel-Gawad AF, El-Mofty MM, Attia SI. Effect of dimethoate on some haematological parameters of toad *Bufo regularis*. J King Saud Univ 1995; 7: 85-93.
39. Harris JA. Seasonal variation in some hematological characteristics of *Rana pipiens*. Comp Biochem Physiol Part A: Physiol 1972; 43: 975-989.
40. Al-Amoudi WM. Haematological and Biochemical Effects of Metalaxyl Fungicide on Albino Mice. Amer J Biochem 2012; 2: 62-66.
41. Nussey G, Van Vuren JHJ, Du Preez HH. Effects of Copper on Haematology and Osmoregulation of the Mozambique Tilapia, *Oreochromis mossambicus* (Cichlidae). Comp Biochem Physiol 1995;111: 369-380.
42. Anand Kumar A, Tripathy AP, Tripathy AK. Effect of dimecron on the blood parameter of *Heteropneustes fossilis*. J Environ Biol 2001; 22: 297-299.
43. Garg VK, Garg SK, Tyagi SK. Manganese induced haematological and biochemical abnormalities in *Heteropneustes fossilis*. J Environ Biol 1989;10: 349-353.
44. Dhanapakiam P, Ramasamy VK. Toxic effect of zinc and copper mixtures on some haematological and biochemical parameters in common carp, *Cyprinus carpio*. J Environ Biol 2001; 22:105-111.
45. Wepener WM, Vuren JHJ. Effect of chromium at different pH values on the haematology of *Tilapia sparrmanii*. J Comp Biochem Physiol 1992;161: 375-381.
46. Isomaa B, Hagerstrand H, Paatero G, Engblom AC. Permeability alterations and anti-hemolysis induced by amphiphiles in human erythrocytes. Biochim Biophys Acta 1986; 869: 510-524.
47. Atamanalp YT, Haliloglu, H, Aras MS. Alterations in the hematological parameters of rainbow trout, *Oncorhynchus mykiss*, exposed to cypermethrin. The Israeli J Aquacult Bamidgeh 2002; 54: 99-103.
48. Parma MJ, Loteste A, Campane M, Baccheta C. Changes in haematological parameters in *Prochilodus lineatus* exposed to sublethal concentration of cypermethrin. J Environ Biol 2007; 28:147-149.
49. Kartheek RM, David M. Assessment of renal toxicity in rats exposed to commercial formulations of fipronil. Int J Pharm Chem Biol Sci 2017; 7:296-303.
50. David M, Kartheek RM. Biochemical changes in liver of freshwater fish *Cyprinus carpio* exposed to sublethal concentration of sodium cyanide. Indo Amer J Pharm Res 2014; 4: 3669-3675.
51. David M, Kartheek RM. In vivo studies on hepato-renal impairments in freshwater fish *Cyprinus carpio* following exposure to sublethal concentrations of sodium cyanide. Environ Sci Pollut Res Int 2016; 23: 722-733.
52. David M, Kartheek RM. Fipronil induced modulations in biochemical and histopathological aspects of male Wistar albino rats: A subchronic study. World J Environ Biosci 2016; 5: 26-32.
53. David M, Kartheek RM. Histopathological alterations in spleen of freshwater fish *Cyprinus carpio* exposed to sublethal concentration of sodium cyanide. Open Vet J 2015; 5:1-5.

Cite this article as:

Kartheek RM, David M. Modulations in Haematological Aspects of Wistar Rats Exposed to Sublethal Doses of Fipronil under Subchronic Duration. J Pharm Chem Biol Sci 2017; 5(3):187-194